

## 2015 Hybrid Winter Rye Regional Trial – Cornell University

Entry	Grain Yield (kg/h)						Test Weight kg/hl	Rank	Lodging Score 0-9	Head Date 2 Loc	Winter Surv. %	Height cm	
	Regional Locations												
	lth-Hel	lth-Ket	MonCo	GenCo	Mean	Rank							
1	Brasetto (180 k/m <sup>2</sup> )	6784	6718	9635	5045	7046	1	71.3	11	2.1	5/26	97	118
2	Brasetto (200 k/m <sup>2</sup> )	6978	6145	8303	4138	6391	13	71.1	13	2.2	5/26	94	121
3	Brasetto (250 k/m <sup>2</sup> )	6535	5843	8498	4241	6279	16	71.2	12	2.8	5/26	88	117
4	KWS Bono (H 119)	6575	6466	8897	4459	6599	7	72.3	6	3.6	5/26	100	116
5	KWS Rhavo (H 120)	6466	6410	8725	4265	6467	10	72.4	4	2.2	5/26	99	115
6	Medina (wheat ck)	4656	3087	5092	ND	4278	**	73.4	2	2.5	5/31	98	92
7	KWS H-139 (Nikko)	6611	6364	8106	4231	6328	15	71.9	8	3.2	5/26	97	119
8	KWS H-140 (Daniello)	6808	6597	8178	4139	6431	11	71.7	9	1.9	5/26	98	118
9	KWS H-141 Livado)	6881	6361	8823	4619	6671	5	72.4	5	2.7	5/26	100	119
10	KWS H-144 (Gatano)	6839	6359	7774	4693	6416	12	71.0	15	2.1	5/26	100	111
11	Danko	5293	5058	7600	2774	5181	17	73.3	3	2.2	5/25	99	135
12	AC Hazlet	4413	4270	6706	2615	4501	18	72.2	7	2.2	5/26	99	136
13	Guttino	6470	6428	8660	4834	6598	8	71.3	10	2.8	5/26	99	116
14	KWS H-145	7312	6665	9277	4021	6819	4	68.4	20	1.4	5/27	99	120
15	KWS H-146	7331	6217	9420	4620	6897	3	69.9	17	1.4	5/27	98	121
16	KWS H-147	7369	4834	8588	4641	6358	14	69.7	18	2.6	5/26	100	116
17	KWS H-149	6490	6348	9673	3950	6615	6	69.3	19	1.0	5/26	100	119
18	KWS H-150	6707	6306	8709	4578	6575	9	71.0	14	1.0	5/26	99	118
19	KWS H-151	7095	6643	9052	4879	6917	2	70.0	16	2.0	5/26	99	114
20	Erie (wheat ck)	5428	3892	5486	ND	4936	**	75.5	1	1.0	6/1	93	82
	Mean	6452	5851	8260	4264	6215		71.5		2.1	5/25	98	116
	CV	4.7	13.3	8.2	15.4								

\*\*Not ranked because of deer damage in Genesee Co.

Entry	Grain Yield				Test Weight		Lodging	Head Date	Height cm	Winter Surv.	
	3 Year		2 Year		2 Year		0-9				
	kg/h	b/a	kg/h	b/a	kg/hl	lb/b	2 Yr				2 Yr
1	Brasetto (180 k/m <sup>2</sup> )	6719	107	7434	118	69.6	55.7	2.4	5/26	123	97
2	Brasetto (200 k/m <sup>2</sup> )	6556	104	6961	111	69.6	55.7	2.2	5/26	125	96
3	Brasetto (250 k/m <sup>2</sup> )	6695	107	7137	114	69.7	55.8	2.3	5/26	123	94
4	KWS Bono (H 119)	6539	104	6994	111	70.8	56.6	3.0	5/26	120	99
5	KWS Rhavo (H 120)	6515	104	6868	109	70.9	56.7	1.9	5/26	122	98
6	Medina (wheat ck)	3997	59	4286	64	71.2	57.0	2.2	5/31	94	99
7	KWS H-139 (Nikko)			6887	110	70.2	56.1	2.8	5/27	123	98
8	KWS H-140 (Daniello)			7027	112	69.9	55.9	2.2	5/26	121	97
9	KWS H-141 Livado)			6930	110	70.6	56.4	2.2	5/27	122	98
10	KWS H-144 (Gatano)			6922	110	69.5	55.6	2.1	5/27	116	99
11	Danko			5791	92	71.8	57.4	2.6	5/25	140	100
12	AC Hazlet			5058	81	70.1	56.1	2.6	5/27	144	100

**M.E. Sorrells, D. Benschler, J. Shiffer, J. Tanaka - Department of Plant Breeding & Genetics, Cornell University**